



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**I. Part of a Letter to the Publisher from Dr Davies  
concerning an unusual Colic.**

Birmingham, Oct. 3d. 1701.

S I R,

**I** Aft *May*, my Ingenious Friend Dr *Shaw* of *Tamworth*, was pleased to desire my assistance in the following case. A person, aged between 50 and 60, had been for 3 or 4 years last past troubled with Gripes, which generally return'd once a month, or thereabouts; his Body for the most part costive, and therefore was forced to spur Nature with *Daffy's* Elixir or Aloes; sometimes the smoaking a Pipe of Tobacco, supply'd the use of these Medicines. Last *April* coming from *London* heated himself very much with his Journey, having walked a good part of the way, and as soon as he came home, had a return of his Colic pains, which continued upon him for 18 days, notwithstanding the methods commonly used in that case, during which time he had no Stool, besides what the first and second Clysters brought away; his complaint upon his seizure was of a pain in his right side in the *Regio Iliaca*. Some time before he dyed his Belly swelled much, and was as tense as any Drum, he vomited for 2 or 3 days at the beginning, which left him, and returned not till just before he dyed, which was at the expiration of the 18th day, at which time he brought up 2 or 3 mouthfuls of black Choler; but never during his whole illness vomited any excrement. Dr *Shaw* who lived upon the spot, prevail'd to have an hasty Inspection of his Abdomen, the Relations (it being a thing *Malæ famæ* in that Neighbourhood) desired it might be done with all secrecy and expedition,  
which

which prevented the Doctors taking so nice a view of all the parts within the Abdomen as he would otherwise have done. He found some black Choler in the Stomach, the *Duodenum* and the rest of the *Intestina tenuia* void of excrement, but incredibly distemper'd with Wind, and tracing the Canalis of the Guts as far as the *Cæcum*, found that of a blackish colour; and from thence for about a yard in length, the Colon mortified and so rotten, that the Excrements had made their way thro' it at several places, into the Cavity of the Abdomen; about 2 inches of the mortified Gut was fastened to the *Peritonæum* on the right side. This part of the Colon was so extended with excrements of a soft consistence, that they when taken out filled two Chamber-pots; at the extremity of the mortification, towards the *Rectum* the obstruction which occasioned all these mistortunes offered it self to view very plainly; for about ten inches of the Colon was doubled, as if you had taken a piece of Tape, and folded it the two contiguous surfaces of the duplicature adhered so firmly together, that you could not separate them without tearing the exterior coat of the Intestine. Upon separating this coalescence, there fell from that part a whitish Mucus, the adhesion was about 3 inches broad; the middle of the Duplicature which made the acute angle, and where the excrements stopped, was smaller, and the Membranes thinner than in any other part of the Gut; from thence towards the *Rectum* the Colon was sound, and void of excrements, occasioned by the frequent use of Clysters.